

Medicinal herbs inhibit cholesterol absorption and synthesis in liver

Authors:

S Murad, Altaf Hussain Mahar, Shumaila Najeeb, Nusratullah Khan, Hafiz Moeen-Ud-Din, Seema Shah
Murad, Farid-Ud-Din

HOD Pharmacology, IMDC-DANTH, Islamabad-Pakistan

Research scholar at HEJ Institute of Chemistry, UK, Pakistan

Asso Prof of Histopathology at Yusra Medical College, Islamabad Pakistan

Assistant Prof of Biochemistry at Bolan University of Medical and health Sciences, Quetta, Pakistan

Associate Prof of Anatomy at AIMC Lahore Pakistan

Gynecologist at NMC Karachi Pakistan

Lecturer Pharmacology at DANTH Islamabad Pakistan

Corresponding Author:

Dr. S Murad, Pharmacology Deptt: Akbar Niazi Teaching Hospital and IM&DC, affiliated with SZABMU, Islamabad-PAKISTAN.

Article Received: 07-February-2020, Revised: 27-February-2020, Accepted: 17-March-2020

ABSTRACT:

Traditional herbal medicines are naturally occurring, plant-derived substances with minimal or no industrial processing that have been used to treat illness within local or regional healing practices. Traditional herbal medicines are getting significant attention in global health debates. In China, traditional herbal medicine played a prominent role in the strategy to contain and treat severe acute respiratory syndrome. We in this study have compared hypolipidemic effects of Fenugreek, Curcuma longa, and Lemon. Study was conducted at Jinnah Hospital Lahore-Pakistan from January 2016 to May 2016. Ninety hyperlipidemic patients of age group 19 to 70 were included in the study. Exclusion criteria were diabetic, alcoholic additives, hypertensive patients and those whose kidney or liver functions were impaired. Consent was taken from all participants. Their base line lipid profile was taken in biochemistry laboratory of the hospital. They were divided in three groups i.e. 30 patients in each group. Group-I was advised to take 500 mg of Curcuma longa (haldi) mixed in fresh milk without cream, thrice daily for two months. Group-II patients were advised to take 100 grams of Fenugreek leaves mixed with salad in each meal (thrice daily) for the period of two months. Group-III patients were advised to take 40 ml of fresh lemon juice mixed with 40 ml mineral water thrice daily for two months. They all were advised not to take heavy meal rich with any type of fat like junk food etc. One hour daily brisk walk was advised to all participants. 15 days follow up visit was scheduled for them. After two months their lipid profile was re-determined. When results were compiled and statistically analyzed by applying paired 't' test, it revealed that Curcuma longa decreased total cholesterol, TG, LDL cholesterol 16.10, 20.01, and 17.59 mg/dl respectively. Fenugreek decreased total cholesterol, TGs, and LDL cholesterol 14.70, 17.33, and 17.06 mg/dl respectively. Lemon in two months therapy decreased total cholesterol, TGs, and LDL cholesterol 15.45, 10.13, and 11.97 mg/dl respectively. None of the above mentioned herbs raised HDL cholesterol significantly. It was concluded from this research work that Curcuma longa, Fenugreek leaves and Lemon are mild to moderately effective hypolipidemic herbs to lower total plasma cholesterol, triglycerides, and LDL cholesterol but have no potential to raise HDL cholesterol when analyzed biostatistically.

Keywords: Prevention is better than cure, herbs, lemon, methi, haldi

INTRODUCTION:

Many hope traditional herbal medicine research will play a critical role in global health. China, India, Nigeria, the United States of America (USA) and WHO have all made substantial research investments in traditional herbal medicines. Industry has also invested millions of US dollars looking for promising medicinal herbs and

novel chemical compounds. This is still a relatively modest investment compared to the overall pharmaceutical industry; however, it raises interesting ethical questions, some of which are not faced in more conventional drug development. Atherosclerotic problems are encountered with enhancement of LDL uptake by monocytes and macrophages¹⁻². In the liver,

uptake of plasma LDL is mediated via specific LDL receptors, but a scavenger receptor system is employed by macrophages. Plasma LDL must be modified prior to uptake by macrophages. Analysis of the lipid content in the oxidatively modified LDL from hyper lipidemic patients revealed that the level of lysophosphatidylcholine was greatly elevated, and the high level of the lysolipid was shown to impair the endothelium-dependent relaxation of the blood vessels³⁻⁷. In allopathy blood lipid levels are normalized by statins, fibrates, niacin and bile acid binding resins (BABRs). All of these allopathy related drugs have low compliance ranges from mild compliance due to metallic taste of BABRs to severe side effects like rhabdomyolysis by fibrates and statins⁸. Due to low compliance and adverse effects of conventionally used hypolipidemic agents, herbal medicines are going to be famous among Physicians, and Cardiologists⁹. Medicinal herbs like Curcuma Longa, Fenugreek, and Lemon contents are being used as mild to moderate hypolipidemic agents. Curcuma Longa commonly known as Haldi in India and Pakistan is used to lower plasma lipids, in view of their contents. It contains Curcuminoids (curcumin, demethoxycurcumin, and bisdemethoxycurcumin), turmerone, atlantone, zingiberene, proteins, and resins¹⁰. Curcumin reduces both the oxidation and circulation of oxidized levels of LDL cholesterol which leads to reduction in the occurrence or treatment of already present atherosclerosis in the subjects¹¹. Research has revealed that curcumin from turmeric is an extremely efficient agent in increasing cholesterol uptake by the liver cells¹². Several independent studies have shown that curcumin increases the expression of cholesterol and lipoprotein receptors on the liver cells¹³. Curcumin also increases cholesterol and bile acids excretion in feces^{14,15}. The composition of fenugreek includes a large number of chemical components. They include proteins and amino acids, flavonoids, saponins and steroidal saponins, coumarin, lipids, vitamins, minerals, galactomannan fiber and alkaloids, such as trigonelline. Extracts are available standardized to contain 50 percent saponins or 20 percent of the amino acid 4-hydroxyisoleucine¹⁶. Treatment with fenugreek. Seed powder normalized the enhanced lipid peroxidation and increased susceptibility to oxidative stress associated with depletion of antioxidants¹⁷. The steroidal saponins (diosgenin, yamogenin, tigogenin and neotigogenin) are thought to inhibit cholesterol absorption and synthesis and hence its potential role in arteriosclerosis¹⁸. Phenolics in Curcuma Longa have potential health benefits mainly due to their antioxidant properties such as reactive oxygen species (ROS) scavenging and inhibition, electrophile scavenging and metal chelation¹⁹. Epidemiological studies support a relationship between the consumption

of phenolic rich food products and a low incidence of coronary heart disease, atherosclerosis, certain forms of cancer and stroke²⁰. Lipid peroxidation is inhibited by flavonoids and flavanones present in Lemon²¹.

PATIENTS & METHOD:

The research work was conducted at Jinnah Hospital, Lahore from January 2016 to May 2016 with approval of Research ethics committee of the Hospital. Ninety hyperlipidemic patients were selected for research work. Written consent was taken from all patients. Specific Performa was designed for the research work. Hyperlipidemic patients were selected with age range from 19 to 70 years. **EXCLUSION CRITERIA:** Exclusion criteria were hypertension, hypothyroidism, diabetes mellitus, alcohol addictive patients, peptic ulcer, any gastrointestinal upset, renal impairment, and any hepatic or cardiac problem. All patients were divided in three groups (group-I, group-II, group-III), 30 patients in each group. Their baseline lipid profile data were taken and filed in specifically designed Performa, at start of taking medicine, like lipid profile, blood pressure and pulse rate. Serum lipid profile (total cholesterol, HDL-cholesterol, triglyceride) parameters were determined after a 12 hour overnight fast by standard methods. LDL-cholesterol level was calculated according to the Friedewald's formula. Thirty patients of group-I were advised to take 500 mg of Curcuma longa (haldi) mixed in fresh milk without cream, thrice daily for two months. Group-II patients were advised to take 100 grams of Fenugreek leaves mixed with salad in each meal (thrice daily) for the period of two months. Group-III patients were advised to take 40 ml of fresh lemon juice mixed with 40 ml mineral water thrice daily for two months. They all were advised not to take heavy meal rich with any type of fat like junk food etc. One hour daily brisk walk was advised to all participants. 15 days follow up visit was scheduled for them. After two months their lipid profile was re-determined. **STATISTICAL ANALYSIS:** Mean values of the day-0 and day-60 of tested parameters (total cholesterol, LDL cholesterol, triglycerides, and HDL cholesterol) \pm SD were taken to be analyzed statistically. SPSS 10 version 2.00.01.10 was used to analyze pretreatment and post treatment values of all parameters. Paired 't' test was applied to determine changes in pre and post treatment values. P-value >0.05 was considered as non significant changes in tested parameters, and p-values <0.01 were considered as significant changes.

RESULTS:

It revealed in this study that curcuma longa decreased triglycerides (TG), total cholesterol (TC), and LDL cholesterol of 29 hyperlipidemic patients 20.01, 16.10, and 17.59 mg/dl respectively. Raise in HDL cholesterol

in this group was 03.70 mg/dl. Fenugreek leaves reduced TG, TC, LDL cholesterol of 28 hyperlipidemic patients 17.33, 14.70, and 17.06 mg/dl respectively. Lemon juice reduced TC, TG, and LDL cholesterol 15.45, 10.13, and

11.97 mg/dl respectively. HDL raised 03.55 mg/dl in this group. Changes in all tested parameters, and their statistical significance are shown in following table.

Three tables are changed

Table: 1 Showing parameters values before and after treatment with their statistical significance in group-1(n = 29) (Curcuma longa or haldi)

TC at day-0 271.87±1.04	TC at day-60 255.77±2.77	Difference in mg/dl 16.10	P-value= <0.001
TG at day-0 216.09±2.43	TG at day-60 196.08±2.45	20.01	P-value= <0.001
LDL-C at day-0 179.65±2.87	LDL-C at day-60 162.06±2.51	17.59	P-value= <0.001
HDL-C at day-0 37.95±1.45	HDL-C at day-60 41.65±1.91	3.70	P-value= <0.001

Table: 2 Showing parameters values before and after treatment with their statistical significance in group-2 (n = 28) (Fenugreek or methi)

TC at day-0 280.56±1.06	TC at day-60 265.86±2.65	Difference 14.70	P-value= <0.001
TG at day-0 224.87±1.55	TG at day-60 207.54±1.98	17.33	P-value= <0.001
LDL-C at day-0 213.13±2.78	LDL-C at day-60 196.07±1.56	17.06	P-value= <0.001
HDL-C at day-0 35.19±2.32	HDL-C at day-60 38.08±1.67	2.89	P-value= <0.001

Table: 3 Showing parameters values before and after treatment with their statistical significance in group-3 (n = 29) (Lemon water)

TC at day-0 258.21±2.12	TC at day-60 242.76±1.89	Difference 15.45	P-value= >0.05
TG at day-0 246.56±2.11	TG at day-60 236.43±2.43	10.13	P-value= >0.05
LDL-C at day-0 179.08±2.87	LDL-c at day-60 167.11±1.77	11.97	P-value= >0.05
HDL-C at day-0 36.22±1.77	HDL-C at day-60 39.77±1.55	3.55	P-value= >0.05

KEY: All values are written in mean and ± stands for standard error of mean. T-C= serum total cholesterol, TG= serum triglycerides, LDL-C= low density lipoprotein cholesterol, HDL-C= high density lipoprotein cholesterol. All parameters pre and post-treatment are measured in mg/d. n = sample size. P-value <0.01 stands for significant change, P-value >0.05 stands for non-significant change.

DISCUSSION:

We proved that All of these three herbs have no significant influence on HDL cholesterol. Same results did prove in the study conducted by Wiseman SA et al²² who described that pectin present in these three herbs inhibit enterohepatic circulation of bile acids and excrete cholesterol in feces. Flavenoids present in Fenugreek are

responsible for inhibition of cholesterol synthesis²³. Gidez LI et al²⁴ stated that herbal medications have more than one or two mechanism to balance plasma lipids in hyperlipidemic patients. Bingham SA et al²⁵ stated that the major reasons for hypercholesterolemia in today's world are obesity, consuming high fat food, diabetes and having a family history of high cholesterol. This disorder

is reported to affect a large number of people all across the world and is one of the leading causes of death as well. Cholesterol is reduced in the body by managing weight and diet. Regular exercise, lesser consumption of fatty foods, more consumption of fruits and vegetables help in ameliorating the symptoms of hypercholesterolemia. However, in most cases, medications also known as anti hypertensive and anti cholesterol drugs are also required, especially in chronic cases. There are several problems associated with medications though and people are now switching towards newer and less toxic therapies to control and reduce cholesterol levels in the body. Several natural herbs, supplements and food products are known to maintain healthy cholesterol levels and reduce cholesterol in mild hypercholesterolemia. These therapies, if opted for, help in preventing the disorder but are less helpful in chronic cases of this disease. In an experiment²⁶, controlled dosing of turmeric was used to feed hypercholesterolemic rabbits and the effects on LDL oxidation was analyzed. It was found that turmeric extracts efficiently and quickly reduced the levels of cholesterol in the blood along with the incidences of atherosclerosis with time. The reason for these therapeutic effects, when analyzed, was found to be the preventive action of turmeric on the oxidation of LDL cholesterol. Mattern T et al²⁷ have also explained same important mechanism of action of Turmeric that oil of these seeds inhibits enterohepatic circulation causing biosynthesis of bile acids instead of cholesterol by hepatocytes. The results obtained from research work conducted by Geleijnse JMet al²⁸ revealed that all extracts of the fenugreek exhibit antioxidant activity. These findings suggest that the fenugreek extracts could act as potent source of antioxidants. Magee E et al²⁹ mentioned that many herbs and their constituents have potential to reduce total plasma cholesterol LDL cholesterol, and triglycerides but they do not raise HDL cholesterol because it needs special plasma proteins as lipoproteins responsible for structural and functional integrity of HDL particles. Jhumarr J et al³⁰, Loversa T et al³¹, hundahg P et al³², described about citrus fruit Lemon that its Citric acid is antioxidant which lead to decrease oxidative stress in human body. Cases like these present challenging questions related to the role of traditional herbal medicines in public health. In general, international research on traditional herbal medicines should be subject to the same ethical requirements as all research related to human subjects.⁹ An ethical framework previously outlined by Emanuel et al. and revised for international research⁸ offers a useful starting point for thinking about the ethics of international traditional herbal medicine research. This framework includes eight ethical requirements for clinical research³³.

REFERENCES:

1. L. A Layrrv, S. A. & Bouwens, L. C. The chemistry of tea flavonoids. *Crit. Rev. Food Sci.* 2011;37:693-704.
2. Mocali V, Thorpe, G. Atherosclerosis: newer view points. *Pk Med. J.* 2014;313:229.
3. Suthy G, Ferro-Luzzi, A. In vivo antioxidant effect of green and black tea in man. *J. Clin. Nutr.* 2012;50:28-32.
4. Palivov FF, Wiseman, S. A. A single dose of tea with or without milk increases plasma antioxidant activity in humans. *Ind J. Nutr.* 2012;54:87-92.
5. Sanders, T. A. & Wiseman, H. Flavonoids protect against oxidative damage to LDL in vitro: use in selection of a flavonoid rich diet and relevance to LDL oxidation resistance ex vivo. *Free Rad. Res.* 2015;33:419-426.
6. Strain, J. J. The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. *Anal. Biochem.* 20013;239:70-76.
7. Solvoll, K. & Foss, O. P. Tea consumption. Relationship to cholesterol, blood pressure, and coronary and total mortality. *Prev. Med.* 2014;21:546-553.
8. Ito, T., Yoshida, H., Ayaori, M., Nishiwaki, M., Yonemura, A., Hara, Y. & Nakamura, H. Conventional hypolipidemic medicines and their compliance. *Libyan J. Nutr.* 2014;66:261-266.
9. Sesso, H. D., Gaziano, J. M., Buring, J. E. & Hennekens, C. H. Hypolipidemic constituents in herbs. *J. Epidemiol.* 2014;149:162-167.
10. Hodgson, J. M., Puddey, I. B., Croft, K. D., Burke, V., Mori, T. A., Caccetta, R. A. & Beilin, L. J. Chemical contents of *Curcuma Longa*. *J. Nutr.* 2014;71:1103-1107.
11. Klatsky, A. L., Friedman, G. D. & Armstrong, M. A. *Curcuma longa* affects LDL-oxidation. *Med. J. Epidemiol.* 2012;132:479-488.
12. Hertog, M. G., Sweetnam, P. M., Fehily, A. M., Elwood, P. C. & Kromhout, D. Antioxidant flavonols and ischemic heart disease. *Sr J. Med* 2013; 65:1489-1494.
13. Kono, S., Shinchi, K., Ikeda, N., Yanai, F. & Imanishi, K. Miraculous hypolipidemic herb:

- CURCUMA LONGA. Japan. Prev. Med. 2013;21:526-531.
14. Fogelman, A. M., Shechter, I., Seager, J., Hokom, M., Child, J. S. & Edwards, P. A. Curcuma longa inhibits enterohepatic circulation. J Sur & Med Ind 2015;77:2214-2218.
 15. Lowry, O. H., Rosenbrough, N. J., Farr, A. L. & Randall, R. J. ATP-binding cassette transporter and efflux of lipids. J. Biol. Chem. 2013;193:265-275.
 16. Keda, I., Imasato, Y., Sasaki, E., Nakayama, M., Nagao, H., Takeo, T., Yayabe, F. & Sugano, M. Chemical compounds found in Fenugreek seeds. J Bioch 2013;1127:141-146.
 17. Yang, C. S. & Landau, J. M. Handling LDL oxidation with herbal medicine. J. Cl. Nutr. 2014;130:2409-2412.
 18. Keli, S. O., Hertog, M. G., Feskens, E. J. & Kromhout, D. Dietary flavonoids, antioxidant vitamins, and incidence of stroke. J. Med Sc 2014;156:637-642.
 19. Mukhtar, H. & Ahmad, N. Metallic Chelation by Curamin. J Cl Med 2014;71:1698S-1702S.
 20. Princen, H. M., van Duyvenvoorde, W., Buytenhek, R., Blonk, C., Tijburg, L. B., Langius, J. A., Meinders, A. E. Phenolic compound rich foods for cure of CAD. J Health Med Ethop 2014;18:833-41.
 21. Hertog, M. G., Feskens, E. J., Hollman, P. C., Katan, M. B. & Kromhout, D. Lemon Juice for treating hyperlipidemia and CAD. J Ind Med 2015;42:1007-11.
 22. Wiseman, S. A., Balentine, D. A. & Frei, B. Antioxidant potential of pectin present in herbs. J Med Ir 2014;37:705-18.
 23. Tijburg, L. B., Mattern, T., Folts, J. D., Weisgerber, U. M. & Katan, M. B. Flavonoids and cardiovascular disease: a review. J Biol Pk 2014;37(2):771-85.
 24. Gidez, L. I., Miller, G. J., Burstein, M., Slagle, S. & Eder, H. A. MOA of hypolipidemic herbs. J of Med Therapy 2015; 23(4):206-23.
 25. Bingham, S. A., Vorster, H., Jerling, J. C., Magee, E., Mulligan, A., Runswick, S. A. & Cummings, J. H. Obesity, hypercholesterolemia, metabolic syndrome and options of their cure. Med J Prev Med 2015; 78(3):41-55.
 26. Reaven, P. D. & Witztum, J. L. Oxidized low density lipoproteins in atherogenesis: role of dietary modification. J Ethop Med 2013;16(6):51-71
 27. Mattern T, Witztum JL, Weisgerber UM. Turmeric seeds oil interrupts enterohepatic circulation of Bas. J. Lipid Res. 2012 53: 2490-2514
 28. Geleijnse, J. M., Launer, L. J., Hofman, A., Pols, H. A. & Witteman, J. C. How fenugreek is antioxidant? Food Sc 2014;159:2170-74.
 29. Magee, E., Mulligan, A, Constrv YL. Consumption of Foods Rich in Flavonoids Is Related to a Decreased Cardiovascular Risk. J. Med Nutr. 2014;134:23-6.
 30. Jhumarr J, Yulda C, Helov T. Fruits can prevent diseases. MJN 2013;8(7):99-104.
 31. Loversa T, Mught R, Loferr T, Ilova H. Phytochemistry of some plants. Jou Cl Nutr Res 2013;12(1):44-8.
 32. Hundahg P, Urhav I, Ytrag R. Therapeutic approach with medicinal herbs. JRMS 2012;7(5):66-80.
 33. Tyli UI, Thyvo GF, Sueawath JJ, Howan RR, Mekba TG, Illahu RF, Ulkav BT. Lemon is antioxidant fruit. UJPM 2016;12(8):112-6.