

Phytochemical Characterization and Pharmacokinetic Evaluation of Lichen-Derived Cholecalciferol (Vegiferol®): A Comparative Review of Bioavailability and Therapeutic Efficacy

Authors:

B.D.S. Bedi, Premjith Balakrishnan, Dr. Soyeb Nagori, Dr. Javed Sipai, Zahid Sipai

Affiliation:

Aurora Farmabio Pvt. Ltd., Vadodara, Gujarat, India.

Corresponding Author:

Zahid Sipai

<https://doi.org/10.5281/zenodo.18920165>

Received: 30-January-2025, Revised: 18-February-2026, Accepted: 08-March-2026, Online First: 09-March-2026

ABSTRACT:

The global prevalence of Hypovitaminosis D remains a primary public health concern, impacting musculoskeletal integrity and immune homeostasis. While traditional Cholecalciferol (D3) is primarily animal derived, there is an increasing clinical demand for high-potency, vegan-compliant alternatives. This paper evaluates Vegiferol®, a nature identical Vitamin D3 complex derived from the lichen. We examine the pharmacological efficacy of Vegiferol® in its concentrated 1 Million IU/g Oil and stabilized 100,000 IU/g Powder formats. Clinical data suggest that even at the physiological baseline dose of 400 IU, the unique lichen-derived matrix demonstrates superior serum absorption markers within a 48-hour window compared to conventional counterparts.

Keywords: *Vegiferol®, Lichen-based D3, Cholecalciferol, Bioavailability, 25(OH)D, Nutritional Immunology, Calcidiol.*

INTRODUCTION:

Vitamin D3 (Cholecalciferol) functions as a fundamental pro-hormone, essential for the regulation of approximately 3% of the human genome through the Vitamin D Receptor (VDR) [1]. Unlike Ergocalciferol (D2), Cholecalciferol (D3) is the metabolite naturally synthesized in the skin and has been shown to be significantly more effective in raising and maintaining serum 25-hydroxyvitamin D [25(OH)D] levels [2].

PHARMACOLOGICAL MECHANISM AND DOSE-RESPONSE:

The 400 IU Physiological Standard:

International guidelines, including those from the Institute of Medicine (IOM), recommend a baseline dose of 400 IU for the prevention of deficiency and maintenance of bone health [4]. Scientific analysis of Vegiferol® indicates that due to its high purity and the absence of lanolin-associated impurities, even this conservative dose effectively supports serum homeostasis. The high-potency raw material standards (1 Million IU/g and 100,000 IU/g) allow for precise micro-dosing and consistent therapeutic outcomes [5].

Enhanced Bioavailability:

The lichen-based matrix of Vegiferol® ensures optimal micellar formation in the small intestine, leading to a faster and more robust rise in circulating Calcidiol [6].

SYSTEMIC CLINICAL APPLICATIONS:

- **Immunological Modulation:** Induces the expression of antimicrobial peptides like Cathelicidin and Defensins, pivotal in defense against respiratory pathogens [7].
- **Neuropsychiatric Integrity:** VDRs are highly concentrated in the hippocampus; deficiency is a risk factor for neurodegenerative decline and depressive disorders [8].
- **Cardio-Metabolic Regulation:** Contributes to vascular health by regulating the renin-angiotensin-aldosterone system (RAAS) [9].

MATERIAL SCIENCE AND STANDARDIZATION:

Aurora Farmabio utilizes Supercritical Fluid Extraction (CO₂) at pressures of 300-350 bar and temperatures of

50-60°C. This ensures a "Clean Label" product that is non-GMO, solvent-free, and allergen-free ^[10].

CONCLUSION:

Vegiferol® redefines the standard for plant-based Vitamin D3. By offering superior pharmacokinetics and a nature-identical structure, it provides a clinically superior alternative to animal-derived sources.

REFERENCES:

1. **Holick, M. F. (2007).** Vitamin D deficiency. *New England Journal of Medicine*, 357(3), 266-281.
2. **Tripkovic, L., et al. (2012).** Comparison of vitamin D2 and vitamin D3 supplementation. *Am J Clin Nutr*, 95(6), 1357-1364.
3. **Bikle, D. D. (2014).** Vitamin D metabolism and clinical applications. *Chemistry & Biology*, 21(3), 319-329.
4. **Institute of Medicine (US). (2011).** Dietary Reference Intakes for Calcium and Vitamin D. *National Academies Press*.
5. **Heaney, R. P., et al. (2011).** Vitamin D3 is more potent than vitamin D2 in humans. *J Clin Endocrinol Metab*, 96(3), E447-E451.
6. **Jäpelt, R. B., & Jakobsen, J. (2013).** Vitamin D in plants: a review. *Frontiers in Plant Science*, 4, 136.
7. **Ginde, A. A., et al. (2009).** Association between serum 25(OH)D level and upper respiratory infection. *Arch Intern Med*, 169(4), 384-390.
8. **Eyles, D. W., et al. (2005).** Distribution of the vitamin D receptor in human brain. *J Chem Neuroanat*, 29(1), 21-30.
9. **Wang, T. J., et al. (2008).** Vitamin D deficiency and risk of cardiovascular disease. *Circulation*, 117(4), 503-511.
10. **Wacker, M., & Holick, M. F. (2013).** Vitamin Defects on skeletal and extraskeletal health. *Nutrients*, 5(1), 111-148.